

1 FILES HAVE ONE OR MORE ANSWERS, 66 FILES SEARCHED IN STNINDEX

L38 QUE L33 AND L34

=> d rank L38

'FULL' IS NOT VALID IN THE CURRENT FILE

This option is not valid in the current file. Enter the command without the option at the arrow prompt (=>). Or, first enter the file in which the saved item created. Then enter the command and option at an arrow prompt in the file.

=> d rank

F1 1 DGENE

=> file f1

COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
24.75	87.01

FULL ESTIMATED COST

FILE 'DGENE' ENTERED AT 14:25:13 ON 14 MAY 2003
COPYRIGHT (C) 2003 THOMSON DERWENT

FILE LAST UPDATED: 9 MAY 2003 <20030509/UP>

DGENE CURRENTLY CONTAINS 3,563,592 BIOSEQUENCES

>>> DOWNLOAD THE DGENE WORKSHOP MANUAL:

http://www.stn-international.de/training_center/bioseq/dgene_wm.pdf

>>> DOWNLOAD COMPLETE DGENE HELP AS PDF:

http://www.stn-international.de/training_center/bioseq/dgene_help.pdf <<<

>>> DOWNLOAD DGENE BLAST/GETSIM FREQUENTLY ASKED QUESTIONS:

<http://www.stn-international.de/service/faq/dgenefaq.pdf> <<<

KAILASH C. SRIVASTAVA

DEX 'ADISCTI, ADISINSIGHT, ADISNEWS, AGRICOLA, ANABSTR, AQUASCI, BIOBUSINESS, BIOCOMMERCE, BIOSIS, BIOTECHABS, BIOTECHDS, BIOTECHNO, CABA, CANCERLIT, CAPLUS, CEABA-VTB, CEN, CIN, CONFSCI, CROPB, CROPU, DDFB, DDFU, DGENE, DRUGB, DRUGLAUNCH, DRUGMONOG2, ...' ENTERED AT 17:06:25 ON 14 MAY 2003

19 FILES HAVE ONE OR MORE ANSWERS
QUE 3-ISOPYLMALATE DEHYDRATASE OR BETA-ISOPROPYLMALATE DEHYDRATASE ORISOP
OPYLMALATE ISOMERASE OR 3-ISOPROPYLMALATE HYDRO-LYASE OR ALPHA-ISOPROP
YLMALATE ISOMERASE

61 FILES HAVE ONE OR MORE ANSWERS
QUE INHIBIT? (5N) (COMPOUND OR MATTER OR COMPD OR SUBSTANCE OR MATERIAL OR
CHEMICAL)

66 FILES HAVE ONE OR MORE ANSWERS
QUE SCREEN? OR EVALUAT? OR ASSAY? OR TEST? OR QUANTI?

60 FILES HAVE ONE OR MORE ANSWERS
QUE (ANTIBIOTIC OR ANTIBACTERIAL OR ANTIFUNGAL OR ANTIMICROBIAL OR FUNGICI
D? OR FUNGISTATIC) (5N) COMPOUND

0 FILES HAVE ONE OR MORE ANSWERS
QUE L1 AND L2

13 FILES HAVE ONE OR MORE ANSWERS,
QUE (EC 4.2.1.33)

10 FILES HAVE ONE OR MORE ANSWERS,
QUE L1 AND L6

0 FILES HAVE ONE OR MORE ANSWERS
QUE L5 AND L7

2 FILES HAVE ONE OR MORE ANSWERS
QUE L7 AND L3

> d rank

1 1 MEDLINE
2 1 USPATFULL

> file f1-f2

> L9

10 2 L9

11 0 L10 AND L2

12 0 L10 AND L4

13 0 L7 AND L4

10 ANSWER 1 OF 2 MEDLINE

N 94171070 MEDLINE

N 94171070 PubMed ID: 8125330

I The LEU1 gene of Ustilago maydis.

U Rubin B P; Li D; Holloman W K

S Department of Microbiology, Cornell University Medical College, New York,
NY 10021.

C GM42482 (NIGMS)

GM42548 (NIGMS)

O GENE, (1994 Mar 11): 140 (1) 131-5.

Journal code: 7706761. ISSN: 0378-1119.

Y Netherlands

T Journal; Article; (JOURNAL ARTICLE)

A English

S Priority Journals

S GENBANK-L20832

M 199404

D Entered STN: 19940420

Last Updated on STN: 19940420

Entered Medline: 19940414

B The nucleotide sequence of the Ustilago maydis LEU1 gene has been
determined. It contains a continuous open reading frame predicted to
encode a protein of 773 amino acids with a molecular mass of 83,234 Da.
The protein is homologous to ***alpha*** - ***isopropylmalate***
isomerases from prokaryotes and eukaryotes, as well as to other
members of a family of structurally related isomerases.

10 ANSWER 2 OF 2 USPATFULL

N 2003:78516 USPATFULL

I STAPHYLOCOCCUS AUREUS POLYNUCLEOTIDES AND SEQUENCES

QH 442.643

IN KUNSCH, CHARLES A., GAITHERSBURG, MD, UNITED STATES
CHOI, GIL A., ROCKVILLE, MD, UNITED STATES
BARASH, STEVEN C., ROCKVILLE, MD, UNITED STATES
DILLON, PATRICK J., GAITHERSBURG, MD, UNITED STATES
FANNON, MICHAEL R., SILVER SPRING, MD, UNITED STATES
ROSEN, CRAIG A., LAYTONSVILLE, MD, UNITED STATES
PI US 2003054436 A1 20030320
AI US 1997-781986 A1 19970103 (8)
PRAI US 1996-9861P 19960105 (60)
DT Utility
FS APPLICATION
LREP HUMAN GENOME SCIENCES INC, 9410 KEY WEST AVENUE, ROCKVILLE, MD, 20850
CLMN Number of Claims: 29
ECL Exemplary Claim: 1
DRWN 2 Drawing Page(s)
LN.CNT 13414

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB The present invention provides polynucleotide sequences of the genome of Staphylococcus aureus, polypeptide sequences encoded by the polynucleotide sequences, corresponding polynucleotides and polypeptides, vectors and hosts comprising the polynucleotides, and ***assays*** and other uses thereof. The present invention further provides polynucleotide and polypeptide sequence information stored on computer readable media, and computer-based systems and methods which facilitate its use.

L14 0 (ALPHA-ISOPROPYLMALATE ISOMERASES) (5N) INHIBIT?
L15 0 (ALPHA-ISOPROPYLMALATE ISOMERASES) (5N) INHIBITOR
L16 0 (ALPHA-ISOPROPYLMALATE ISOMERASES) AND L4

L6 ANSWER 68 OF 70 BIOTECHDS COPYRIGHT 2003 THOMSON DERWENT AND
ISI

AN 1991-08933 BIOTECHDS

TI Some properties of the leucine-biosynthesizing enzymes from *Candida maltosa*;

1-isopropylmalate-synthase, 3-isopropylmalate-dehydratase,
3-isopropylmalate-dehydrogenase purification and characterization

AU Bode R; Birnbaum D

LO Ernst-Moritz-Arndt-Universitaet Greifswald, Fachrichtung Biologie,
Institut fuer Biochemie, Jahnstrasse 15a, O-2200 Greifswald, Germany.

SO J.Basic Microbiol.; (1991) 31, 1, 21-26

CODEN: JBMIEQ

DT Journal

LA English

AN 1991-08933 BIOTECHDS

AB The enzymes of the leucine biosynthetic pathway in *Candida maltosa* L4
were partially purified and their catalytic properties determined.

Maximum activity of the first enzyme, alpha-isopropylmalate
(IPM)-synthase (1-isopropylmalate-synthase, EC-4.1.3.12), was observed at
pH values between 7.5 and 8.8. The K_m values for alpha-ketoisovalerate
and acetyl-CoA were 0.57 mM and 0.064 mM, respectively. Enzyme activity
was ***inhibited*** specifically by L-leucine, and was strongly
dependent on the presence of monovalent cations, preferably K^+ (80 mM).

IPM-dehydratase (***3*** - ***isopropylmalate*** -

dehydratase , EC-4.2.1.33) activity showed a sharp optimum at pH
8.5. The enzyme did not require cations for activity, and L-leucine,
L-isoleucine and L-valine did not ***inhibit*** activity. The pH
optimum of beta-IPM-dehydrogenase (3-isopropylmalate-dehydrogenase,
EC-1.1.1.85) was 6.8, with 50% of the optimum activity expressed at pH
5.8 and pH 7.4. Monovalent cations were not required for dehydrogenase
activity, but divalent ions increased activity (preferably Mn^{2+} at 1 mM).
Enzyme activity was ***inhibited*** by L-valine. (23 ref

IUBMB Enzyme Nomenclature

EC 4.2.1.33

Common name: 3-isopropylmalate dehydratase

Reaction: 3-isopropylmalate = 2-isopropylmaleate + H₂O

Other name(s): β-Isopropylmalate dehydratase; isopropylmalate isomerase; α-isopropylmalate isomerase

Systematic name: 3-isopropylmalate hydro-lyase

Comments: The enzyme also hydrates the product to 3-hydroxy-4-methyl-3-carboxypentanoate, thus bringing about an interconversion between the two isomers.

Links to other databases: [BRENDA](#), [EXPASY](#), [KEGG](#), [WIT](#), CAS registry number: 37290-72-5

References:

1. Gross, S.R., Burns, R.O. and Umbarger, H.E. The biosynthesis of leucine. II. The enzymic isomerization of β-carboxy-β-hydroxyisocaproate and α-hydroxy-β-carboxyisocaproate. *Biochemistry* 2 (1963) 1046-1052.

[EC 4.2.1.33 created 1972, modified 1976]

[Return to EC 4.2.1 home page](#)

[Return to EC 4.3 home page](#)

[Return to EC 4 home page](#)

[Return to Enzymes home page](#)

[Return to IUBMB Biochemical Nomenclature home page](#)

